

## OCTAGAM 5 and 10% (ADULT)

**Use IBW in all adult patients except those whose actual body weight is less than IBW and obese patients greater than 100 kg or BMI greater than 30 kg/m<sup>2</sup> (use adjusted)**

400 MG/KG	
Weight	Dose (g)
50	20
55	20
60	25
65	25
70	30
75	30
80	30
85	35
90	35
95	40
100	40
105	40
110	45
115	45
120	50
125	50
130	50
135	50
140	55
145	60
150	60

500 MG/KG	
Weight	Dose (g)
50	25
55	25
60	30
65	30
70	35
75	35
80	40
85	40
90	45
95	45
100	50
105	50
110	55
115	55
120	60
125	60
130	65
135	65
140	70
145	70
150	75

1 G/KG	
Weight	Dose (g)
50	50
55	55
60	60
65	65
70	70
75	75
80	80
85	85
90	90
95	95
100	100
105	105
110	110
115	115
120	120
125	125
130	130
135	135
140	140
145	145
150	150

2 G/KG	
Weight	Dose (g)
50	100
55	110
60	120
65	130
70	140
75	150
80	160
85	170
90	180
95	190
100	200
105	210
110	220
115	230
120	240
125	250
130	260
135	270
140	280
145	290
150	300

Round doses to the nearest 5 gm in adults (smaller vials reserved for pediatrics)  
**Octagam 5% is the preferred product** and available in 1g, 2.5g, 5g, 10g vials  
 Octagam 10% is restricted for selected patients (i.e. fluid overload) and is available in: 2g, 5g, 10g, 20g vials

**OCTAGAM 5% (ADULT) Supplemental Dose Rounding Table (Preferred Product)**

<b>100 MG/KG</b>	
<b>Weight</b>	<b>Dose (g)</b>
50	5
55	5.5
60	6
65	6.5
70	7
75	7.5
80	8
85	8.5
90	9
95	9.5
100	10
105	11
110	11
115	12
120	12.5
125	13
130	13
135	14
140	14
145	15
150	15

<b>150 MG/KG</b>	
<b>Weight</b>	<b>Dose (g)</b>
50	7.5
55	8.5
60	9
65	10
70	11
75	11
80	12
85	12.5
90	14
95	14
100	15
105	16
110	17
115	17.5
120	18
125	19
130	30
135	20
140	21
145	22
150	22.5

Round doses to the nearest vial size in adults utilizing all vial sizes  
 Octagam 5% vial size: 1g, 2.5g, 5g, 10g

Use IBW in all adult patients except those whose actual body weight is less than IBW and obese patients greater than 100 kg or BMI greater than 30 kg/m<sup>2</sup> (use adjusted)

**OCTAGAM 10% (ADULT) Supplemental Dose Rounding Table**

<b>100 MG/KG</b>	
<b>Weight</b>	<b>Dose (g)</b>
50	5
55	6
60	6
65	6
70	7
75	7
80	8
85	8
90	9
95	9
100	10
105	10
110	11
115	11
120	12
125	12
130	13
135	13
140	14
145	15
150	15

<b>150 MG/KG</b>	
<b>Weight</b>	<b>Dose (g)</b>
50	7
55	8
60	9
65	10
70	10
75	11
80	12
85	13
90	13
95	14
100	15
105	16
110	16
115	17
120	18
125	18
130	20
135	20
140	21
145	22
150	22

Round doses to the nearest vial size in adults by utilizing all vial sizes  
 Octagam 10% is restricted to select patients (i.e. fluid overload) and is available in: 2g, 5g, 10g, 20g  
 Use IBW in all adult patients except those whose actual body weight is less than IBW and obese patients greater than 100 kg or BMI greater than 30 kg/m<sup>2</sup> (use adjusted)